



A novel role for RIP1 kinase in mediating TNF α production

Citation

Christofferson, D E, Y Li, J Hitomi, W Zhou, C Upperman, H Zhu, S A Gerber, S Gygi, and J Yuan. 2012. A novel role for RIP1 kinase in mediating TNF α production. *Cell Death & Disease* 3(6): e320.

Published Version

doi:10.1038/cddis.2012.64

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10459027>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

A novel role for RIP1 kinase in mediating TNF α production

DE Christofferson¹, Y Li¹, J Hitomi¹, W Zhou¹, C Upperman¹, H Zhu¹, SA Gerber¹, S Gygi¹ and J Yuan^{*1}

Receptor-interacting protein 1 (RIP1) is a Ser/Thr kinase with both kinase-dependent and kinase-independent roles in death receptor signaling. The kinase activity of RIP1 is required for necroptosis, a caspase-independent pathway of programmed cell death. In some cell types, the inhibition of caspases leads to autocrine production of TNF α , which then activates necroptosis. Here, we describe a novel role for RIP1 kinase in regulating TNF α production after caspase inhibition. Caspase inhibitors activate RIP1 kinase and another protein, EDD, to mediate JNK signaling, which stimulates Sp1-dependent transcription of TNF α . This pathway is independent of nuclear factor κ B and also occurs after Smac mimetic/IAP antagonist treatment or the loss of TNF receptor-associated factor 2 (Traf2). These findings implicate cIAP1/2 and Traf2 as negative regulators of this RIP1 kinase-dependent TNF α production pathway and suggest a novel role for RIP1 kinase in mediating TNF α production under certain conditions.

Cell Death and Disease (2012) 3, e320; doi:10.1038/cddis.2012.64; published online 14 June 2012

Subject Category: Immunity

Receptor-interacting protein 1 (RIP1) is a multi-functional signal transducer involved in mediating nuclear factor κ B (NF- κ B) activation, apoptosis, and necroptosis. The kinase activity of RIP1 is critically involved in mediating necroptosis, a caspase-independent pathway of necrotic cell death.^{1,2} RIP1 kinase and necroptosis are activated by death receptor ligands, such as TNF α and Fas, when apoptosis is blocked.^{1,3} Necrostatin-1 (Nec-1), a small molecule inhibitor of RIP1 kinase activity, can also block necroptosis. In some cell types, such as mouse fibrosarcoma L929 cells, necroptosis is activated by inhibition of caspase activity using a pan-caspase inhibitor such as zVAD.fmk.³ How RIP1 kinase is activated to mediate necroptosis induced by caspase inhibition is not clear.

RIP1 is ubiquitinated by the cellular inhibitor of apoptosis proteins, cIAP1 and cIAP2.^{4,5} RIP1 is ubiquitinated by a number of other E3 ubiquitin ligases as well, suggesting that RIP1 ubiquitination might regulate RIP1 activity. Smac mimetics (SMs) are a class of compounds modeled after the N terminus of a cellular protein, Smac/DIABLO, that inhibits the IAPs. SMs are under development as anti-cancer drugs.^{6,7} In some cell types, SM treatment can induce autocrine TNF α production and cell death, although the pathway has not been fully elucidated.^{7–10}

TNF α is an important pro-inflammatory cytokine involved in mediating cell death and inflammation in many human diseases such as rheumatoid arthritis and cancers. In a genome-wide siRNA screen to identify genes involved in necroptosis, we found that knockdown of *tnfr1* or treatment with a TNF α -neutralizing antibody was protective against zVAD.fmk-induced cell death in L929 cells, indicating that

zVAD.fmk was likely inducing autocrine TNF α production.¹¹ As knockdown of RIP1 protects against zVAD.fmk-induced death,¹¹ we tested the hypothesis that RIP1 might act upstream of TNF α production after zVAD.fmk and identified a novel function of RIP1 kinase in mediating TNF α production.

EDD/UBR5/hHYD is a putative tumor suppressor and HECT (homologous to E6-AP C-terminus)-domain-containing E3 ubiquitin ligase implicated in cellular pathways including the regulation of gene expression, the DNA damage response, and in necroptosis after it was identified in a siRNA screen.^{11,12} EDD regulates gene expression transcriptionally, by forming complexes with transcription factors, and translationally by regulating protein levels of Paip2, a poly-A-binding protein inhibitor.^{13,14} EDD is also important in the cellular DNA damage response, mediating ATM phosphorylation of its substrates CHK2 and p53 after DNA damage to control cell cycle arrest.^{15–17} Given its multiple functions in mediating cellular processes, EDD likely acts as a chaperone protein, coordinating the various protein complexes involved in different cellular pathways.

In this study, we describe a novel RIP1 kinase-dependent TNF α production pathway occurring in cellular models of necroptosis and apoptosis. We explored this novel TNF α production pathway using a combination of chemical inhibitors and genetic analysis, and defined a protein complex containing EDD, RIP1, and cIAP1 that mediates JNK activation and transcription of TNF α . This TNF α production pathway requires RIP1 kinase and is activated specifically in response to zVAD.fmk stimulation, SM compounds, or TNF receptor-associated factor 2 (Traf2) deficiency.

¹Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA

*Corresponding author: J Yuan, Department of Cell Biology, Harvard Medical School, 240 Longwood Avenue, SGM 400, Boston, MA 02115, USA. Tel: 617 432 4170; Fax: 617 432 4177; E-mail: jyuan@hms.harvard.edu

Keywords: RIP1; TNF α ; necroptosis; necrostatin; inflammation

Abbreviations: RIP1, receptor-interacting protein 1; NF- κ B, nuclear factor κ B; NLS, nuclear localization signal; Traf2, TNF receptor-associated factor 2; Nec-1, necrostatin-1; SMs, Smac mimetics; RSV, Rous sarcoma virus; SV40 LT, SV40 large T antigen

Received 03.2.12; revised 17.4.12; accepted 18.4.12; Edited by H-U Simon

Results

RIP1 and EDD are required for TNF α production in response to zVAD.fmk. To directly examine whether zVAD.fmk stimulates the production of TNF α , we measured TNF α levels after zVAD.fmk treatment. TNF α could be detected in dying L929 cells treated with zVAD.fmk. Nec-1, a RIP1 kinase inhibitor, blocked the increase in TNF α protein levels as well as cell death (Figures 1a and b).

As the knockdown of RIP1 can have differential effects from the inhibition of its kinase activity alone,¹¹ we used Nec-1 as a tool to examine the role of RIP1 kinase in TNF α production. Although Nec-1 has been shown to be a specific inhibitor of RIP1 kinase,² we further tested the specificity of Nec-1 to ensure its suitability for this study. Nec-1 specifically binds RIP1 with K_d 5.6 nM for racemic Nec-1 and K_d 3.1 nM for *R*-Nec-1 (Supplementary Figure S1a). Using KINOMEScan (Ambit Biosciences, San Diego, CA, USA),¹⁸ Nec-1 (10 μ M) was tested for activity against 485 kinases and activated mutant kinases. When ranked in order of inhibition, RIP1 is the top kinase inhibited by Nec-1 (Supplementary Figure S1b). Aside from RIP1, no kinases were inhibited greater than 60% by Nec-1. PDGFR β was inhibited by 72%, however, the K_d of binding of Nec-1 to PDGFR β was greater than 30 μ M, suggesting a false positive (Supplementary Figures S1a and b). Nec-1 is more specific than 35 known kinase inhibitors including imatinib (Gleevec) (Supplementary Figure S1c). As further evidence, Nec-1 can only protect against necroptosis in *Rip1*^{+/+} MEF cells, but not in *Rip1*^{-/-} MEFs (Supplementary Figure S1d). Thus, we conclude that Nec-1 is a highly specific inhibitor of RIP1 kinase activity and an appropriate tool with which to study the specific role of RIP1 kinase.

L929 cells are exquisitely sensitive to death induced by TNF α treatment, so to directly study the effect of RIP1 on TNF α production we tested different cell types that produce TNF α in response to zVAD.fmk treatment. A mouse macrophage cell line, J774, was found to produce easily measurable TNF α levels in response to zVAD.fmk stimulation (Figure 1c). Both primary macrophages and macrophage cell lines undergo necroptosis in response to zVAD.fmk.^{11,19} Although J774 cell treatment with zVAD.fmk induced necroptosis, cell death was not dependent on the production of TNF α . Cell death was observed beginning at least 12 h after the increase in TNF α was first detected and neutralization of TNF α was not sufficient to block zVAD.fmk-induced necroptosis of J774 cells (Figure 1b). Inhibition of RIP1 kinase by Nec-1 completely blocked the production of TNF α in J774 cells, suggesting that RIP1 kinase is required for TNF α production in zVAD.fmk-treated J774 cells (Figure 1c).

The production of TNF α in response to zVAD.fmk treatment was blocked by CHX suggesting that *de novo* protein synthesis is involved (Figure 1d). Consistent with this possibility, our siRNA screen found a significant enrichment of transcription factors and nucleic acid-binding genes among hits protecting against zVAD.fmk-induced necroptosis.¹¹ Treatment with zVAD.fmk activates *de novo* synthesis of TNF α through a mechanism dependent upon the kinase activity of RIP1.

To find additional components of the RIP1 kinase-dependent pathway of TNF α production, we identified RIP1-binding proteins. 293T cells were transfected with a vector expressing Flag-tagged RIP1 kinase, and RIP1 immunocomplexes were isolated using anti-Flag. The binding proteins were identified by mass spectrometry analysis. This analysis identified FADD, a known RIP1-binding protein, thus validating the




Figure 1 RIP1 kinase activates TNF α production. (a) TNF α levels determined by TNF α ELISA and normalized to total protein in lysate from L929 cells treated with 20 μ M zVAD.fmk \pm 10 μ M Nec-1 for 16 h. (b) L929 or J774 cells treated with 20 μ M zVAD.fmk \pm 1.0 μ g/ml anti-TNF α or 10 μ M Nec-1. Cell viability measured by ATP assay after 24 h. (c) TNF α levels in J774 cells treated with 20 μ M zVAD.fmk \pm 10 μ M Nec-1 for 4 and 8 h were determined as in a. (d) TNF α levels in J774 cells treated with 20 μ M zVAD.fmk \pm 0.5 μ g/ml CHX. * P < 0.05, data shown are mean \pm S.E.M., n = 3

experiment. To distinguish binding proteins that have a functional role in mediating the production of TNF α , we compared the list of mass spectrometry-identified proteins with the hits identified in the genome-wide siRNA screen for genes involved in mediating necroptosis of L929 cells in response to zVAD.fmk.¹¹ EDD, encoded by the gene *edd1*, is both a RIP1-binding protein and a gene whose knockdown blocks zVAD.fmk-induced necroptosis. We confirmed that RIP1 coimmunoprecipitated EDD (Figure 2a). Interestingly, overexpression of EDD consistently increases the levels of exogenously expressed RIP1 protein but has no effect on endogenous RIP1, suggesting that overexpressed RIP1 is stabilized by EDD (Figure 2a, Supplementary Figure S2a). Knockdown of EDD protected against zVAD.fmk but not TNF α -induced necroptosis (Figure 2b), indicating EDD does not have any role in TNF α -induced necroptosis and instead has a likely role in zVAD.fmk-induced TNF α production. To determine whether EDD is required for TNF α production, we generated stable knockdown cell lines using retroviral infection of an shRNA construct against EDD. Knockdown of EDD inhibited zVAD.fmk-induced TNF α , demonstrating that EDD is required for TNF α production (Figure 2c).

To determine how TNF α production is activated, we used real-time PCR to measure TNF α mRNA levels after zVAD.fmk treatment. Stimulation with zVAD.fmk increased TNF α mRNA, and treatment with Nec-1, or knockdown of EDD, was able to block the increase in TNF α mRNA (Figure 2d). Thus, RIP1 and EDD activate the transcription of TNF α after zVAD.fmk treatment.

SM induces TNF α in a manner dependent on RIP1 kinase and EDD. To explore the physiological relevance of this

pathway, we looked for other stimuli that could also activate this RIP1 kinase-dependent pathway of TNF α production. SM induces autocrine TNF α production in some cells.^{7–10} Thus, we tested whether RIP1 and EDD have role in SM-induced TNF α . Treatment with SM induces the auto-ubiquitination and degradation of cIAP1 and cIAP2, and also blocks XIAP binding to and inhibition of caspases. We tested the effect of the previously described SM-164^{20,21} on L929 cells and found that, similar to zVAD.fmk, SM-164 induced TNF α -dependent necrotic cell death that could be inhibited by Nec-1 or a TNF α -neutralizing antibody (Figure 3a). Knockdown of RIP1, EDD, or TNFR1 was able to block SM-induced necroptosis, suggesting that SM might be activating the same RIP1- and EDD-dependent pathway of TNF α production as zVAD.fmk (Figure 3b). ELISA was used to detect an increase in TNF α in the lysate of SM-treated L929 cells. SM-induced TNF α could be inhibited by Nec-1 treatment and in EDD-knockdown cells, confirming that SM activates RIP1 kinase and EDD-dependent TNF α production (Figure 3c).

Traf2, an E3 ubiquitin ligase, is constitutively bound to cIAP1/2 within the cell.^{22,23} Traf2-knockout mice have elevated levels of serum TNF α ,²⁴ which suggested to us that Traf2 might act similarly to cIAP1/2 to inhibit induction of TNF α . Indeed, knockdown of Traf2 in L929 cells induced TNF α -dependent necroptosis that could be blocked by both TNF α -neutralizing antibody or knockdown of TNFR1 (Figure 3d). Similar to SM- and zVAD.fmk-induced cell death, Traf2 knockdown-induced death could be blocked by Nec-1 and by knockdown of RIP1 or EDD (Figure 3d). Thus, the absence of Traf2, similar to the loss of cIAP1/2 during SM treatment, activates RIP1- and EDD-dependent TNF α production.




Figure 2 EDD mediates TNF α production. **(a)** 293T cells transfected with Flag-EDD and Flag-RIP1 as indicated. Cell lysates and anti-EDD immunoprecipitates were western blotted with anti-RIP1 and anti-EDD to show both endogenous and overexpressed protein. **(b)** L929 cells transfected for 48 h with siRNA targeting RIP1 or one of the four different siRNAs targeting EDD were treated for 24 h with 20 μ M zVAD.fmk or 10 ng/ml hTNF α and cell viability was measured by ATP assay (top) or lysates collected for western blot (bottom) to check protein knockdown. **(c)** J774 cells (left) or L929 cells (right) stably expressing shRNA against EDD or an empty vector control were treated with 20 μ M zVAD.fmk and TNF α measured as in **a**. EDD knockdown shown in Figure 6e and Supplementary Figure S3a. **(d)** Real-time PCR of RNA isolated from J774 cells treated with zVAD.fmk with or without Nec-1 (left) or EDD stable knockdown J774 cells treated with zVAD.fmk (right). Relative levels of TNF α transcript were determined compared with GAPDH (left) or 18S rRNA (right) and the fold change was calculated by comparing with DMSO-treated cells. * P < 0.05, data shown are mean \pm S.E.M., for all experiments n = 3




Figure 3 Traf2 and cIAP1 inhibit RIP1- and EDD-dependent TNF α production. (a) ATP assay of L929 cells treated with 100 nM SM-164 \pm TNF α -neutralizing antibody or 10 μ M Nec-1 for 24 h. (b) Viability of L929 cells transfected with siRNA against RIP1, EDD, TNFR1, or negative control (NC) for 48 h and treated with 20 μ M zVAD.fmk or 100 nM SM-164 for 24 h. (c) TNF α levels were measured by ELISA in L929 cells treated with 100 nM SM-164 \pm 10 μ M Nec-1 (left) or stably expressing EDD shRNA or vector control (pSRP) and treated with 100 nM SM-164 (right). (d) ATP assay of L929 cells transfected with siRNA against Traf2 and treated for 72 h, with 1.0 μ g/ml anti-mTNF α antibody or 10 μ M Nec-1 (left). For western blot, the cells were treated with 10 μ M Nec-1 from the start of transfection to block cell death and harvested after 48 h. L929 cells transfected with siRNAs targeting Traf2 and either RIP1, EDD, or TNFR1. NC \pm siRNA treated with zVAD.fmk used as positive control. Viability was calculated as the ratio of the luminescence values of Traf2 + siRNA to untreated NC + siRNA (right). * P < 0.05, data shown are mean \pm S.E.M., for all experiments n = 3

SM-induced TNF α production has been studied in human cancer cell lines including breast, ovarian, and lung cancer cells. To validate the role of this RIP1- and EDD-dependent pathway in a previously established model of autocrine TNF α production, we tested the effect of Nec-1 and EDD knockdown on SM-induced apoptosis in the human breast cancer MDA-MB-231 cells. MDA-MB-231 cells undergo TNF α -dependent apoptosis with 100 nM SM-164.²⁰ As has been previously shown, knockdown of RIP1 is able to block SM-induced apoptosis (Figure 4a).⁷ However, neither Nec-1 treatment nor knockdown of EDD inhibited 100 nM SM-induced apoptosis (Figures 4a and b). Surprisingly, we found that a 1000-fold lower dose of SM-164 (0.1 nM) is sufficient to induce TNF α -dependent apoptosis, with death blocked by a TNF α -neutralizing antibody or by caspase inhibition. This 'low dose' of SM induces RIP1 kinase-dependent cell

death that is blocked by Nec-1 or by knockdown of EDD (Figure 4c).

To determine whether RIP1 kinase induces TNF α transcription in the MDA-MB-231 cell model, we measured TNF α mRNA levels by real-time PCR after treatment with either 0.1 nM or 100 nM SM-164. Both doses of SM increased TNF α transcription, but Nec-1 was only able to inhibit TNF α transcription in the cells stimulated with the low dose of SM (Figure 4d). The high, 100 nM dose of SM might be activating additional or non-specific pathways that contribute to TNF α transcription and cell death in a RIP1 kinase-independent manner. Low-dose SM activates a specific RIP1 kinase-dependent mechanism of TNF α transcription.

A complex of EDD, RIP1, and the E3 ligase cIAP1 regulates TNF α production. To confirm the complex of




Figure 4 Smac mimetic induces RIP1 kinase- and EDD-dependent TNF α production in MDA-MB-231 cells. (a) ATP assay of MDA-MB-231 cells transfected with NC, RIP1, or EDD siRNA for 48 h and treated with 100 nM SM-164 for 24 h. (b) Time course of MDA-MB-231 cells treated with 100 nM SM-164 \pm 20 μ M zVAD.fmk or 10 μ M Nec-1. Cell viability was measured by ATP assay. (c) ATP assay of MDA-MB-231 cells treated with 0.1 nM SM-164 \pm 20 μ M zVAD.fmk, anti-hTNF α -neutralizing antibody, 10 μ g/ml CHX, or 10 μ M Nec-1 for 24 h (left). MDA-MB-231 cells transfected with the indicated siRNA for 48 h and treated with 0.1 nM SM-164 for 24 h or harvested for western blot (right). (d) RT-PCR of RNA from MDA-MB-231 cells treated with 0.1 nM SM-164 (top) or 100 nM SM-164 (bottom) \pm 10 μ M Nec-1 for 2, 4, or 8 h as indicated. Relative levels of TNF α transcript determined by normalizing to 18S rRNA transcript and fold change calculated in comparison with DMSO-treated cells. * P < 0.05, data shown are mean \pm S.E.M., for all experiments n = 3, western blots were repeated two times with similar results

EDD with RIP1, we examined the interaction of EDD and RIP1. We found that RIP1 coimmunoprecipitates with endogenous EDD in a constitutive manner and this binding is unaffected by RIP1 kinase activation with zVAD.fmk stimulation (Figure 5a) or after SM treatment (Figure 5b). Additionally, cIAP1, the target of SM, also coimmunoprecipitates with EDD unaffected by zVAD.fmk treatment (Figure 5c).

To understand the interactions between the proteins in this complex, we determined the binding domains of RIP1 and cIAP1 to EDD. We tested the ability of EDD to bind RIP1 truncation mutants lacking the kinase domain (Δ KD), the death domain (Δ DD), and both the intermediate and death domains (Δ C). Each of these RIP1 truncation mutants expressed similarly, except for RIP1 Δ KD, which expressed as several bands. This is not due to loss of its kinase activity as the kinase inactive RIP1 K45M does not show this expression pattern (Supplementary Figure S2b). Each of the RIP1 truncation constructs was able to coimmunoprecipitate with EDD, however, the amount of Δ KD in the immunocomplex was enriched compared with that of Δ C or Δ DD, suggesting that EDD predominantly interacts with the non-kinase domain of RIP1 (Figure 5d).

EDD was expressed with cIAP1 or a related family member, XIAP, to confirm the specificity of binding. Antibodies against EDD coimmunoprecipitated cIAP1, but not XIAP (Figure 5e). Testing the binding of cIAP1 deletion constructs indicated that

BIR1 and BIR2 of cIAP1 are the minimal regions to coimmunoprecipitate with EDD, and are pulled down with the same efficiency as full-length cIAP1, suggesting that BIR1-BIR2 is sufficient for cIAP1 to bind EDD (Figure 5f). There was no binding observed of the BIR3, CARD, or RING-containing constructs with EDD, indicating these domains do not interact with EDD. The BIR domains of cIAP1 also mediate its interactions with other proteins such as Traf2, Smac, and RIP1.^{25,26} EDD may act as a scaffold and bind to both cIAP1 and RIP1 kinase. This complex is important for regulating downstream signaling and the activation of TNF α transcription after zVAD.fmk or SM treatment.

A NF- κ B independent mode of TNF α production. Both the canonical and non-canonical NF- κ B pathways (NF- κ B1 and NF- κ B2, respectively) are activated by SM treatment; inhibition of NF- κ B1 blocks TNF α production and cell death.^{9,10} Loss of either cIAP1/2 or Traf2 stabilizes NIK, inducing the proteasomal processing and activation of NF- κ B2.^{9,10,27} Indeed, NF- κ B2 is processed and activated within hours of SM-164 treatment in both 100 nM SM-164-treated MDA-MB-231 cells and SM-treated L929 cells (Supplementary Figure S2c, Figure 6a). Low-dose SM (0.1 nM) induced NF- κ B2 processing in MDA-MB-231 cells as well, albeit with slower kinetics, likely due to the slower rate of cIAP1 degradation (Supplementary Figure S2c). Importantly, Nec-1 had no effect on NF- κ B2 processing in




Figure 5 Complex of EDD, RIP1, and cIAP1 activate TNF α production. (a) J774 cells treated with 20 μ M zVAD.fmk \pm 10 μ M Nec-1 and immunoprecipitated with anti-EDD or an isotype-matched control antibody (C). The immunoprecipitated proteins were western blotted with anti-RIP1 (representative of three independent experiments). (b) L929 cells were treated with 100 nM SM-164 and EDD immunoprecipitated from cells and western blotted for RIP1. (c) L929 cells treated with 20 μ M zVAD.fmk or 100 nM SM-164 for 7 h and EDD immunoprecipitated and western blotted for cIAP1 interaction. (d) EDD immunoprecipitated from 293T cells expressing EDD and the indicated Flag-RIP1 construct. IP was western blotted with anti-Flag to detect RIP1 interaction. (e) 293T cells expressing EDD and Flag-clAP1, Flag-XIAP, or vector control. EDD immunoprecipitated and western blotted with anti-Flag to detect IAP binding. (f) EDD immunoprecipitated from 293T cells expressing indicated Flag-clAP1 construct and western blotted with anti-Flag to detect interaction of clAP1 construct. All experiments were repeated at least two times. * Indicates a non-specific band

either low-dose SM-treated MDA-MB-231 cells or SM-stimulated L929 cells, indicating that RIP1 kinase does not mediate NF- κ B2 activation (Figure 6a, Supplementary Figure S2c). We were unable to detect the degradation of the inhibitor protein I κ B α , a marker for NF- κ B1 activation, under any stimulation (Figure 6a, Supplementary Figures S2c and d). As NF- κ B is not activated in a RIP1 kinase-dependent manner after either zVAD.fmk or SM stimulation, NF- κ B is

unlikely to have a role in cell death or TNF α production as induced by zVAD.fmk or SM in L929 and J774 cells.

Although we did not observe RIP1 kinase-dependent activation of NF- κ B, to verify that NF- κ B does not have a role in zVAD.fmk- or SM-induced cell death, we knocked down NF- κ B1 and NF- κ B2 and tested the effect on cell death. Consistent with previously published work,²⁸ knockdown of NF- κ B1 and NF- κ B2 were not protective against necroptosis




Figure 6 RIP1 and EDD activate JNK signaling and TNF α production. (a) L929 cells treated with 20 μ M zVAD.fmk or 100 nM SM-164 \pm 10 μ M Nec-1 for 2, 4, and 8 h. Cells lysates were western blotted to determine NF- κ B activation (representative of two independent experiments). (b) WT J774 cells (top) or stable EDD knockdown cells (bottom) treated with 1 μ g/ml LPS \pm Nec-1 for 2 h and TNF α levels measured by ELISA in the cell lysate (NS). (c) ATP assay of L929 cells treated as indicated \pm 10 μ M Nec-1 or 1 μ g/ml JNK inhibitor SP600125 for 24 h. (d) J774 TNF α mRNA levels determined by RT-PCR after treatment with 20 μ M zVAD.fmk \pm 1 μ g/ml SP600125 for 2 and 4 h. Fold change calculated by comparing with GAPDH control and normalizing to untreated cells. (e) J774 cells (left) or J774 EDD stable knockdown cell lines (right) were treated with zVAD.fmk \pm Nec-1 as indicated for 15 min, 30 min, 1, 2, or 4 h and western blotted with anti-phospho-JNK (pJNK) (representative of three independent experiments). (f) ATP assay of L929 cells transfected with siRNA targeting Sp1 for 48 h and treated with 20 μ M zVAD.fmk, 100 nM SM-164, or 10 ng/ml hTNF α for 24 h. Cell lysates western blotted to confirm knockdown of Sp1. * P < 0.05, data shown are mean \pm S.E.M., for all experiments n = 3

(Supplementary Figure S2e). An inhibitor of NF- κ B, SN50,²⁹ did not block zVAD.fmk-induced TNF α release in J774 cells (Supplementary Figure S2f). Thus, we conclude that TNF α production induced by zVAD.fmk or SM is independent of NF- κ B activity.

Lipopolysaccharide (LPS) is a pro-inflammatory stimulus found on the outer membrane of bacteria that activates TNF α transcription. Although LPS-induced TNF α is dependent on NF- κ B, we tested the role of RIP1 and EDD in this pathway of TNF α production. Neither Nec-1 nor EDD knockdown was

able to block LPS-induced TNF α (Figure 6b). Thus, RIP1 and EDD activation of TNF α production is distinct from NF- κ B-dependent pathways such as LPS-induced TNF α .

JNK signaling activates TNF α transcription downstream of RIP1 and EDD. Our siRNA screen identified a number of transcription factors as hits, including c-Jun and Sp1, both of which can be activated by JNK and MAPK signaling.^{11,30,31} The importance of c-Jun/AP-1 in mediating TNF α transcription and cell death in response to zVAD.fmk has already been confirmed.^{19,28} Inhibition of JNK signaling blocks TNF α transcription induced by zVAD.fmk (Figure 6d). Consistently, increased phosphorylation of JNK, indicating JNK kinase activation, was observed shortly after zVAD.fmk treatment. A definitive role for RIP1 kinase in activating JNK signaling has not been previously shown. We found that JNK phosphorylation after zVAD.fmk was attenuated in Nec-1-treated and EDD-knockdown cells (Figure 6e, Supplementary Figure S3a). SM stimulation also activates JNK signaling in a manner dependent on RIP1 kinase and EDD (Supplementary Figure S3b). Thus, RIP1 kinase and EDD activate JNK signaling to induce TNF α transcription.

TNF α itself can also activate JNK. Nec-1, however, has no effect on TNF α -induced JNK phosphorylation (Supplementary Figure S3c). AIP1/Dab2IP, an ASK1 (JNK MAP3K) interacting protein, is reportedly a substrate of RIP1 after TNF α stimulation.³² However, neither knockdown of AIP1/Dab2IP nor knockdown of ASK1 in our siRNA screen¹¹ protected against zVAD.fmk-induced cell death (Supplementary Figure S3d). TNF α -induced JNK activation is RIP1 kinase independent and is activated by a separate pathway from zVAD.fmk treatment.

Analysis of our siRNA screen showed a significant enrichment of screen hits with Sp1-binding sites in their promoters, suggesting that Sp1 regulates transcription of genes required for necroptosis.¹¹ Consistent with this possibility, we found that knockdown of Sp1 specifically inhibits zVAD.fmk- or SM-induced cell death but not TNF α -induced necroptosis, similar to EDD knockdown (Figure 6f). Sp1 is able to synergize with c-Jun/AP-1 to activate TNF α transcription.³³ It is likely that both of these transcription factors are activated by JNK signaling downstream of RIP1 kinase and EDD to promote transcription of TNF α and possibly other genes in response to zVAD.fmk or SM stimulation.

Discussion

RIP1 kinase has been previously shown to specifically mediate TNF α -induced necroptosis downstream of TNFR1 by regulating the formation of complex IIb.^{34,35} In this study, we demonstrate a novel function of RIP1 kinase involving its interaction with EDD to regulate JNK activation and TNF α production. This pathway of TNF α production is activated specifically in response to treatment with zVAD.fmk or SM, or by knockdown of Traf2 and is distinguishable from the TNF α production pathway regulated by NF- κ B that can be activated by TLR signaling (Figure 7). We show that activation of this




Figure 7 Model of RIP1- and EDD-mediated pathway of TNF α production. RIP1 kinase-dependent TNF α production can be activated by the treatment with zVAD.fmk, SM to degrade cIAP1/2, or by knockdown of Traf2. EDD, RIP1, and cIAP1 are constitutively bound and RIP1 kinase and EDD activate JNK signaling after stimulation. Traf2 and cIAP1 also constitutively interact. JNK activates the transcription factors AP-1 and Sp1, which transcribe TNF α mRNA. Nec-1 inhibits the activation of JNK and production of TNF α . TNF α produced can elicit inflammatory responses and induce cell death in the surrounding cells

RIP1 kinase-dependent pathway leads to TNF α transcriptional activation. Our examination of this pathway indicates that RIP1 kinase and EDD mediate a common pathway of JNK activation and TNF α production in mouse and human systems, cell types such as macrophages and breast cancer cells, and in cells capable of undergoing either apoptosis or necroptosis. Our study suggests that RIP1 kinase not only regulates necroptosis downstream of TNFR1 signaling, but also has an important role in mediating the production of TNF α .

The role of RIP1 kinase in activating TNF α production is distinct from its role in mediating necroptosis. The function of RIP1 in TNF α production may provide a possible explanation for situations where Nec-1 was found to protect against apoptosis. MDA-MB-231 cells treated with a low dose of SM undergo typical TNF α -dependent apoptosis that can be blocked by Nec-1. This is due Nec-1 inhibiting the production of TNF α , which is required for apoptosis to occur after SM treatment, rather than a role of RIP1 kinase in mediating apoptosis itself. SM are currently in clinical trials as an anti-cancer treatment, indicating the role of RIP1 kinase in mediating the production of TNF α may be relevant for SM activity *in vivo*.

Both cIAP1/2 and Traf2 have been implicated as E3 ubiquitin ligases targeting RIP1.^{5,36} The finding that loss of either cIAP1/2 or Traf2 can activate RIP1 suggests that these proteins normally function to keep RIP1 inactive. *traf2*^{-/-} mice are normal at birth but become progressively runted and die prematurely with elevated serum TNF α levels.²⁴ The lethal phenotype is rescued by the loss of TNF α or TNFR1 in double-knockout *traf2*^{-/-} *tnf*^{-/-} or *traf2*^{-/-} *tnfr1*^{-/-} mice.³⁷ Our study suggests the possibility that RIP1 kinase regulated JNK activation mediates the production of TNF α in these mouse models of human diseases. The identification of SM and Traf2 knockdown as inducers of RIP1 kinase-mediated

TNF α production suggests that RIP1 kinase activation might be regulated directly by ubiquitination by cIAP1/2 and/or Traf2.

SMs, in addition to inducing the degradation of cIAP1 and cIAP2, and inhibit the activity of XIAP.^{6,20} Many of the cellular effects of SM in sensitizing cells to apoptosis have been attributed to the loss of caspase inhibition by XIAP.^{8,20} However, the effect of SM on RIP1 is likely due to the degradation of cIAP1/2, as the effect of SM on TNF α production was recapitulated in cIAP1-knockout cells but not XIAP-deficient cells.¹⁰ Furthermore, EDD specifically interacts with cIAP1, so it is likely that they act in the same pathway.

EDD has been implicated in diverse cellular processes such as the DNA damage response and gene expression. EDD constitutively binds RIP1 kinase and the E3 ubiquitin ligase cIAP1. We propose that EDD functions as a scaffold protein in this pathway and interacts with the critical regulatory proteins. Similar to its role in mediating ATM phosphorylation of its substrates p53 and CHK2,^{15–17} EDD may also mediate RIP1 phosphorylation of its substrate(s) in this pathway. We propose that EDD and RIP1 kinase mediate the activation of JNK signaling, potentially via recruitment of a RIP1 substrate that activates JNK to mediate multiple signaling pathways that are regulated by JNK. Future work is needed to determine how RIP1 kinase and EDD activate the JNK signaling pathway.

Materials and Methods

Gene knockdown experiments. L929 and MDA-MB-231 cells were reverse transfected with 25–50 nM siRNA (Dharmacon, Lafayette, CO, USA) in 384-well plates (Corning, Lowell, MA, USA) using HiPerfect transfection reagent (Qiagen, Gaithersburg, MD, USA), according to the manufacturers protocol. After 48 h of transfection, the cells were treated with zVAD.fmk, hTNF α , or SM-164. After 18–24 h, cell viability was determined by ATP assay using the CellTiter-Glo Luminescent Cell Viability Assay (Promega, Fitchburg, WI, USA). Stable knockdown cell lines were generated using a retroviral or lentiviral expression system to infect L929 or J774 cells, respectively. Expression of pSRP or pLKO.1 and packaging vectors were transfected into 293T cells to generate virus. L929 or J774 cells were infected with the virus and selected in medium containing 10 μ g/ml or 2 μ g/ml puromycin, respectively.

TNF α ELISA. J774 cells were plated 1×10^5 cells/well in 24-well plates or L929 cells plated 4×10^6 cells/10-cm plate, treated, and the cell supernatant collected or cells lysed in 1% Triton X-100 in PBS supplemented with Complete Protease Inhibitor (Roche, Indianapolis, IN, USA). Total protein levels were determined by Bradford protein assay (Bio-rad, Hercules, CA, USA). The levels of TNF α in cell lysate or supernatant were quantified using the mTNF α ELISA kit according to the manufacturers instructions (R&D Systems, Minneapolis, MN, USA).

Real-time PCR. To determine mRNA levels of TNF α in the cell, RNA was harvested according to the manufacturers protocol using the Qiagen RNeasy kit. One microgram of RNA was reverse transcribed with random hexamers and SuperScript II First Strand cDNA Synthesis System (Invitrogen, Carlsbad, CA, USA). The qPCR performed using $2 \times$ SYBR green master mix on the ABI 7900HT qPCR machine (Applied Biosystems, Carlsbad, CA, USA). Fold change in RNA was calculated using the comparative Ct method, normalizing to 18S rRNA or GAPDH control. Taqman probes were used for mTNF α and GAPDH (Applied Biosystems). Primers used are as follows: mTNF α (forward: 5'-CTTCTCAT TCCTGCTTGTGG-3', reverse: 5'-ATGAGAGGGAGGCCATTG-3'), hTNF α (forward: 5'-GAGGCCAAGCCCTGGTATG-3', reverse: 5'-CGGGCCGATTGATCT CAGC-3' PrimerBank ID 25952110b2),³⁸ or 18S rRNA (forward: 5'-CCTCGGGCTT AATTTGACTC-3', reverse: 5'-AGACAAATCGCTCCACCAAC-3').

Coimmunoprecipitation. 293T cells were transfected by the calcium phosphate method and lysed after 24 h in 50 mM Tris-Cl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 10% glycerol, and protease inhibitor cocktail (Roche). L929 cells were lysed in the buffer described except with 50 mM NaCl and 0.5% NP-40. The indicated antibody was used with protein A/G sepharose beads (Thermo Scientific, Waltham, MA, USA) for immunoprecipitation. IP was analyzed by SDS-PAGE and western blot.

Mass spectrometry of RIP1 interacting proteins. 293T cells were transfected by the calcium phosphate method with Flag-tagged RIP1 for 48 h. The cells were lysed (20 mM HEPES (pH 7.3), 5 mM EDTA, 150 mM NaCl, 5 mM NaF, 0.2 mM NaVO₃, 1% Triton X-100, complete protease inhibitor cocktail) and immunoprecipitated using anti-Flag M2 agarose (Sigma, St. Louis, MO, USA). The beads were washed $5 \times$ with lysis buffer and the bound proteins eluted using 150 ng/ μ l Flag peptide. The eluted proteins were TCA precipitated and identified by mass spectrometry.

Expression vectors. The following plasmids were used: pcDNA3-Flag-RIP1 WT, Δ KD, Δ C, and Δ DD. RIP1 Δ KD was constructed using the following primers: 5'-CGAATCCGGAATTCGGCCGACATT-3' and 5'-TGCAGACTCGAGGTTT TGGCTGACGTAAAT-3' to PCR a fragment from nt 843–2007 (NM_003804). RIP1 Δ DD was constructed using the forward primer 5'-ACGATGACGATAAAG AATTCAGGATGCAA-3', and the reverse primer 5'-AGGTGCTCGAGCGTCAG ACTAGTGGTATT-3' to PCR a fragment from nt 1–1751 (NM_003804). RIP1 Δ C was constructed using the same forward primer as the Δ DD construct and the reverse primer 5'-TTCTTCTAATTGCTCGAGATAAAAAGGCCT-3' to PCR a fragment from nt 1–885 (NM_003804). Each of the PCR fragments was cloned into pcDNA3 using EcoRI and XhoI. pCMV-Tag2b-Flag-EDD (courtesy of C.K.W. Watts); Flag-XIAP; Flag-cIAP1, WT and the truncation constructs BIR1-2, BIR1-3, BIR3-RING, CARD-RING, RING. The shRNAs targeting EDD were inserted into the empty vector backbone (sequences: 5'-TGACAGCAGAACATAATT-3' in pSRP; 5'-GCTCGTCTTGATCTACTTTAT-3' in pLKO.1) (courtesy of K.P. Lu).

Antibodies and reagents. The primary antibodies used were anti-RIP1 (BD Biosciences, San Jose, CA, USA), anti-EDD (Novus Biologicals for IP, rat monoclonal raised against aa 459–794 for WB), anti-pJNK, anti-JNK, and anti-NF- κ B2 (Cell Signaling), anti-NF- κ B1, anti-I κ B α , and anti-Traf2 (Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-cIAP1, anti-Flag M2 (Sigma), and anti-mouse and anti-human TNF α -neutralizing antibodies (R&D Systems). The compounds used were zVAD.fmk, hTNF α (Cell Sciences, Canton, MA, USA), SM-164 (kindly provided by Dr. Xiaomeng Wang),^{20,21} 7-Cl-O-Nec-1,³ SP600125 (A.G. Scientific, San Diego, CA, USA), SN50/SN50M (Calbiochem, Darmstadt, Germany), and LPS (Sigma).

Tissue culture. L929, J774, and 293T cells were maintained in DMEM supplemented with 10% FBS and penicillin and streptomycin (Invitrogen). MDA-MB-231 cells were grown in RPMI with 10% FBS, penicillin, and streptomycin (Invitrogen).

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgements. This work was supported in part by a NIH F31 Predoctoral Fellowship (to DEC) and grants from the National Institute of Aging (R37-012859) and a NIH Director's Pioneer Award (to JY). The data in Supplementary Figures S1a–c were kindly provided by TetraLogic Pharmaceuticals, Inc. We thank Drs. Nahum Sonenberg and KP Lu for helpful discussion. We also thank Dr. Xiaomeng Wang for the generous gift of SM-164, Dr. KP Lu for shRNA vectors for EDD, Dr. CKW Watts for the mammalian EDD expression vector, and Dr. Michelle Kelliher for *Rip1* +/+ and *Rip1* –/– MEFs.

- Holler N, Zaru R, Micheau O, Thome M, Attinger A, Valitutti S *et al*. Fas triggers an alternative, caspase-8-independent cell death pathway using the kinase RIP as effector molecule. *Nat Immunol* 2000; 1: 489–495.
- Degterev A, Hitomi J, Gernsheid M, Ch'en IL, Korkina O, Teng X *et al*. Identification of RIP1 kinase as a specific cellular target of necrostatins. *Nat Chem Biol* 2008; 4: 312–321.

3. Degterev A, Huang Z, Boyce M, Li Y, Jagtap P, Mizushima N *et al*. Chemical inhibitor of nonapoptotic cell death with therapeutic potential for ischemic brain injury. *Nat Chem Biol* 2005; **1**: 112–119.
4. Bertrand MJ, Milutinovic S, Dickson KM, Ho WC, Boudreau A, Durkin J *et al*. cIAP1 and cIAP2 facilitate cancer cell survival by functioning as E3 ligases that promote RIP1 ubiquitination. *Mol Cell* 2008; **30**: 689–700.
5. Park SM, Yoon JB, Lee TH. Receptor interacting protein is ubiquitinated by cellular inhibitor of apoptosis proteins (c-IAP1 and c-IAP2) *in vitro*. *FEBS Lett* 2004; **566**: 151–156.
6. Li L, Thomas RM, Suzuki H, De Brabander JK, Wang X, Harran PG. A small molecule Smac mimic potentiates TRAIL- and TNF α -mediated cell death. *Science New York, NY* 2004; **305**: 1471–1474.
7. Petersen SL, Wang L, Yalcin-Chin A, Li L, Peyton M, Minna J *et al*. Autocrine TNF α signaling renders human cancer cells susceptible to Smac-mimetic-induced apoptosis. *Cancer Cell* 2007; **12**: 445–456.
8. Gaither A, Porter D, Yao Y, Borawski J, Yang G, Donovan J *et al*. A Smac mimetic rescue screen reveals roles for inhibitor of apoptosis proteins in tumor necrosis factor- α signaling. *Cancer Res* 2007; **67**: 11493–11498.
9. Varfolomeev E, Blankenship JW, Wayson SM, Fedorova AV, Kayagaki N, Garg P *et al*. IAP antagonists induce autoubiquitination of c-IAPs, NF- κ B activation, and TNF α -dependent apoptosis. *Cell* 2007; **131**: 669–681.
10. Vince JE, Wong WW, Khan N, Feltham R, Chau D, Ahmed AU *et al*. IAP antagonists target cIAP1 to induce TNF α -dependent apoptosis. *Cell* 2007; **131**: 682–693.
11. Hitomi J, Christofferson DE, Ng A, Yao J, Degterev A, Xavier RJ *et al*. Identification of a molecular signaling network that regulates a cellular necrotic cell death pathway. *Cell* 2008; **135**: 1311–1323.
12. Callaghan MJ, Russell AJ, Woollatt E, Sutherland GR, Sutherland RL, Watts CK. Identification of a human HECT family protein with homology to the Drosophila tumor suppressor gene hyperplastic discs. *Oncogene* 1998; **17**: 3479–3491.
13. Henderson MJ, Russell AJ, Hird S, Munoz M, Clancy JL, Lehrbach GM *et al*. EDD, the human hyperplastic discs protein, has a role in progesterone receptor coactivation and potential involvement in DNA damage response. *J Biol Chem* 2002; **277**: 26468–26478.
14. Yoshida M, Yoshida K, Kozlov G, Lim NS, De Crescenzo G, Pang Z *et al*. Poly(A) binding protein (PABP) homeostasis is mediated by the stability of its inhibitor, Paip2. *EMBO J* 2006; **25**: 1934–1944.
15. Henderson MJ, Munoz MA, Saunders DN, Clancy JL, Russell AJ, Williams B *et al*. EDD mediates DNA damage-induced activation of CHK2. *J Biol Chem* 2006; **281**: 39990–40000.
16. Munoz MA, Saunders DN, Henderson MJ, Clancy JL, Russell AJ, Lehrbach G *et al*. The E3 ubiquitin ligase EDD regulates S-phase and G(2)/M DNA damage checkpoints. *Cell cycle Georgetown, Tex* 2007; **6**: 3070–3077.
17. Ling S, Lin WC. EDD inhibits ATM-mediated phosphorylation of p53. *J Biol Chem* 2011; **286**: 14972–14982.
18. Fabian MA, Biggs WH III, Treiber DK, Atteridge CE, Azimioara MD, Benedetti MG *et al*. A small molecule-kinase interaction map for clinical kinase inhibitors. *Nat Biotechnol* 2005; **23**: 329–336.
19. Yu L, Alva A, Su H, Dutt P, Freundt E, Welsh S *et al*. Regulation of an ATG7-beclin 1 program of autophagic cell death by caspase-8. *Science New York, NY* 2004; **304**: 1500–1502.
20. Lu J, Bai L, Sun H, Nikolovska-Coleska Z, McEachern D, Qiu S *et al*. SM-164: a novel, bivalent Smac mimetic that induces apoptosis and tumor regression by concurrent removal of the blockade of cIAP-1/2 and XIAP. *Cancer Res* 2008; **68**: 9384–9393.
21. Sun H, Nikolovska-Coleska Z, Lu J, Meagher JL, Yang CY, Qiu S *et al*. Design, synthesis, and characterization of a potent, nonpeptide, cell-permeable, bivalent Smac mimetic that concurrently targets both the BIR2 and BIR3 domains in XIAP. *J Am Chem Soc* 2007; **129**: 15279–15294.
22. Rothe M, Pan MG, Henzel WJ, Ayres TM, Goeddel DV. The TNFR2-TRAF signaling complex contains two novel proteins related to baculoviral inhibitor of apoptosis proteins. *Cell* 1995; **83**: 1243–1252.
23. Shu HB, Takeuchi M, Goeddel DV. The tumor necrosis factor receptor 2 signal transducers TRAF2 and c-IAP1 are components of the tumor necrosis factor receptor 1 signaling complex. *Proc Natl Acad Sci USA* 1996; **93**: 13973–13978.
24. Yeh WC, Shahinian A, Speiser D, Kraunus J, Billia F, Wakeham A *et al*. Early lethality, functional NF- κ B activation, and increased sensitivity to TNF-induced cell death in TRAF2-deficient mice. *Immunity* 1997; **7**: 715–725.
25. Samuel T, Welsh K, Lober T, Togo SH, Zapata JM, Reed JC. Distinct BIR domains of cIAP1 mediate binding to and ubiquitination of tumor necrosis factor receptor-associated factor 2 and second mitochondrial activator of caspases. *J Biol Chem* 2006; **281**: 1080–1090.
26. Wang L, Du F, Wang X. TNF- α induces two distinct caspase-8 activation pathways. *Cell* 2008; **133**: 693–703.
27. Zarnegar BJ, Wang Y, Mahoney DJ, Dempsey PW, Cheung HH, He J *et al*. Noncanonical NF- κ B activation requires coordinated assembly of a regulatory complex of the adaptors cIAP1, cIAP2, TRAF2 and TRAF3 and the kinase NIK. *Nat Immunol* 2008; **9**: 1371–1378.
28. Wu YT, Tan HL, Huang Q, Sun XJ, Zhu X, Shen HM. zVAD-induced necroptosis in L929 cells depends on autocrine production of TNF α mediated by the PKC-MAPKs-AP-1 pathway. *Cell Death Differ* 2010; **18**: 26–37.
29. Lin YZ, Yao SY, Veach RA, Torgerson TR, Hawiger J. Inhibition of nuclear translocation of transcription factor NF- κ B by a synthetic peptide containing a cell membrane-permeable motif and nuclear localization sequence. *J Biol Chem* 1995; **270**: 14255–14258.
30. Derijard B, Hibi M, Wu IH, Barrett T, Su B, Deng T *et al*. JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* 1994; **76**: 1025–1037.
31. Benasciutti E, Pages G, Kenzior O, Folk W, Blasi F, Crippa MP. MAPK and JNK transduction pathways can phosphorylate Sp1 to activate the uPA minimal promoter element and endogenous gene transcription. *Blood* 2004; **104**: 256–262.
32. Zhang H, Zhang H, Lin Y, Li J, Pober JS, Min W. RIP1-mediated AIP1 phosphorylation at a 14-3-3-binding site is critical for tumor necrosis factor-induced ASK1-JNK/p38 activation. *J Biol Chem* 2007; **282**: 14788–14796.
33. Falvo JV, Uglierolo AM, Brinkman BM, Merika M, Parekh BS, Tsai EY *et al*. Stimulus-specific assembly of enhancer complexes on the tumor necrosis factor α gene promoter. *Mol Cell Biol* 2000; **20**: 2239–2247.
34. Cho YS, Challa S, Moquin D, Genga R, Ray TD, Guildford M *et al*. Phosphorylation-driven assembly of the RIP1-RIP3 complex regulates programmed necrosis and virus-induced inflammation. *Cell* 2009; **137**: 1112–1123.
35. He S, Wang L, Miao L, Wang T, Du F, Zhao L *et al*. Receptor interacting protein kinase-3 determines cellular necrotic response to TNF- α . *Cell* 2009; **137**: 1100–1111.
36. Lee TH, Shank J, Cusson N, Kelliher MA. The kinase activity of Rip1 is not required for tumor necrosis factor- α -induced I κ B kinase or p38 MAP kinase activation or for the ubiquitination of Rip1 by Traf2. *J Biol Chem* 2004; **279**: 33185–33191.
37. Nguyen LT, Duncan GS, Mirtsos C, Ng M, Speiser DE, Shahinian A *et al*. TRAF2 deficiency results in hyperactivity of certain TNFR1 signals and impairment of CD40-mediated responses. *Immunity* 1999; **11**: 379–389.
38. Spandidos A, Wang X, Wang H, Seed B. PrimerBank: a resource of human and mouse PCR primer pairs for gene expression detection and quantification. *Nucleic Acids Res* 2010; **38**, Database issue D792–D799.



Cell Death and Disease is an open-access journal published by Nature Publishing Group. This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on Cell Death and Disease website (<http://www.nature.com/cddis>)